Brain Research, 77 (1996) 231–232
© 1996 Elsevier Science B.V.

Author Index

Bardo, M.T., Donohew, R.L. and Harrington, N.G.
Psychobiology of novelty seeking and drug seeking behavior, 23
Borkowski, J., see Mondadori, C., 223

Cabib, S., see Castellano, C., 1
Cassady, J.M.
Avoidance and classical conditioning of leg flexion in dogs, 79
Castellano, C., Cabib, S. and Puglisi-Allegra, S.
Psychopharmacology of memory modulation: Evidence for multiple interaction among neurotransmitters and hormones, 1
Challet, E., Le Maho, Y., Pévet, P., Nobelis, P. and Malan, A.
Ventromedial hypothalamic lesions prevent the fasting-induced changes in day-night pattern of locomotor activity, 155
Chapouthier, G., see Guillot, P.-V., 211
Coleman-Mesches, K., Salinas, J.A. and McGaugh, J.L.
Unilateral amygdala inactivation after training attenuates memory for reduced reward, 175
Coles, B.L.K., see Whishaw, I.Q., 135

Deacon, R.M.J. and Rawlins, J.N.P.
Effects of aspiration lesions of hippocampus or overlying neocortex on concurrent and configural object discriminations in rats, 165
Decety, J.
The neurophysiological basis of motor imagery, 45
Donohew, R.L., see Bardo, M.T., 23

Eckersdorf, B., Golębiewski, H. and Konopacki, J.
Kainic acid versus carbachol induced emotional-defensive response in the cat, 201

Gibb, R., see Kolb, B., 125
Gispen, W.-H., see van Rijzingen, I.M.S., 149
Golębiewski, H., see Eckersdorf, B., 201
Gorny, G., see Kolb, B., 125
Guillot, P.-V. and Chapouthier, G.
Intermale aggression and dark/light preference in ten inbred mouse strains, 211
Guimarães, F.S., see Netto, S.M., 215

Halpern, M., see Shapiro, L.S., 101
Harrington, N.G., see Bardo, M.T., 23
Hasegawa, T., see Sakai, N., 181

Kameyama, T., see Katoh, A., 219
Katoh, A., Nabeshima, T., Kuno, A., Wada, M., Ukai, R. and Kameyama, T.
Changes in striatal dopamine release in stress-induced conditioned suppression of motility in rats, 219
Kogure, K., see Sakai, N., 181
Kolb, B., Ladowski, R., Gibb, R. and Gorny, G.
Does dendritic growth underly recovery from neonatal occipital lesions in rats, 125
Konopacki, J., see Eckersdorf, B., 201
Kubota, K., see Nakamura, K., 53
Kuno, A., see Katoh, A., 219

Ladowski, R., see Kolb, B., 125
Le Maho, Y., see Challet, E., 155
Li, C.S., see Shapiro, L.S., 101

Malan, A., see Challet, E., 155
Marshall, J.F., see Vargo, J.M., 189
McGaugh, J.L., see Coleman-Mesches, K., 175
Möbius, H.-J., see Mondadori, C., 223
Moebius, H.-J., see Mondadori, C., 227
Mondadori, C., Möbius, H.-J. and Borkowski, J.
The $GABA_B$ receptor antagonist CGP 36 742 and the nootropic oxiracetam facilitate the formation of long-term memory, 223
Mondadori, C., Moebius, H.-J. and Zingg, M.
CGP 36 742, an orally active $GABA_B$ receptor antagonist, facilitates memory in a social recognition test in rats, 227

Nabeshima, T., see Katoh, A., 219
Nagasawa, H., see Sakai, N., 181
Nakamura, K. and Kubota, K.
The primate temporal pole: its putative role in object recognition and memory, 53
Netto, S.M. and Guimarães, F.S.
Role of hippocampal 5-HT_{1A} receptors on elevated plus maze exploration after a single restraint experience, 215
Nobelis, P., see Challet, E., 155

Peretz, B., see Srivatsan, M., 115
Pévet, P., see Challet, E., 155
Puglisi-Allegra, S., see Castellano, C., 1

Rawlins, J.N.P., see Deacon, R.M.J., 165
Roland, R.M., see Shapiro, L.S., 101
Ryu, J.H., see Sakai, N., 181

Sakai, N., Yanai, K., Ryu, J.H., Nagasawa, H., Hasegawa, T., Sasaki, T., Kogure, K. and Watanabe, T.
Behavioral studies on rats with transient cerebral ischemia induced by occlusion of the middle cerebral artery, 181
Salinas, J.A., see Coleman-Mesches, K., 175
Sasaki, T., see Sakai, N., 181
Shapiro, L.S., Roland, R.M., Li, C.S. and Halpern, M.
Vomeronasal system involvement in response to conspecific odors in adult male opossums, *Monodelphis domestica*, 101
Spruijt, B.M., see van Rijzingen, I.M.S., 149
Srivatsan, M. and Peretz, B.
Effect of acetylcholinesterase inhibition on behavior is age-dependent in freely moving *Aplysia*, 115

Ukai, R., see Katoh, A., 219

van Doremalen, E., see van Rijzingen, I.M.S., 149
van Rijzingen, I.M.S., van Doremalen, E., Gispen, W.-H. and Spruijt, B.M.
Long-term impoverished housing effects on morris maze performance after a fimbria lesion, 149
Vargo, J.M. and Marshall, J.F.

Unilateral frontal cortex ablation producing neglect causes time-dependent changes in striatal glutamate receptors, 189

Wada, M., see Katoh, A., 219
Watanabe, T., see Sakai, N., 181
Whishaw, I.Q. and Coles, B.L.K.
Varieties of paw and digit movement during spontaneous food handling in rats: Postures, bimanual coordination, preferences, and the effect of forelimb cortex lesions, 135

Yanai, K., see Sakai, N., 181

Zingg, M., see Mondadori, C., 227